

3000 K Street, N.W., Suite 500  
P.O. Box 25696  
Washington, D.C. 20007-8696  
(202) 672-5300

TO: Assistant Commissioner for Patents  
Box Patent Applications  
Washington D.C. 20231

02/02/99  
  
U.S. PTO

Attorney Docket No. 017283/0123  
(must include alphanumeric codes if no inventors named)

**UTILITY PATENT APPLICATION TRANSMITTAL**  
**(new nonprovisional applications under 37 CFR 1.53(b))**

Transmitted herewith for filing is the patent application of:

**INVENTOR(S):** Patrick BENOIT; Francois MEYER; Deborah MAGUIRE; Ivan PLAVEC; and Michael G. TOVEY

**TITLE:** MONOCLONAL ANTIBODIES AGAINST THE INTERFERON RECEPTOR, WITH NEUTRALIZING ACTIVITY AGAINST TYPE 1 INTERFERON

In connection with this application, the following are enclosed:

**APPLICATION ELEMENTS:**

Specification - 30 TOTAL PAGES

(preferred arrangement:)

- Descriptive Title of the Invention
- Cross Reference to Related Applications
- Statement Regard Fed sponsored R&D
- Reference to Microfiche Appendix
- Background of the Invention
- Brief Summary of the Invention
- Brief Description of the Drawings (if filed)
- Detailed Description
- Claim(s)
- Abstract of the Disclosure

Drawings - Total Sheets 5

Declaration and Power of Attorney - Total Sheets 4

Newly executed (original or copy)

Copy from a prior application (37 CFR 1.63(d))

(relates to continuation/divisional boxes completed) - NOTE: Box below

DELETION OF INVENTOR(S) - Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d) (2) and 1.33(b).

Incorporation By Reference (useable if copy of prior application Declaration being submitted)

The entire disclosure of the prior application, from which a COPY of the oath or declaration is supplied as noted above, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

Microfiche Computer Program (Appendix)

Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)

Computer Readable Copy

Paper Copy (identical to computer copy)

Statement verifying identify of above copies

**ACCOMPANYING APPLICATION PARTS**

Assignment Papers (cover sheet & document(s))  
 37 CFR 3.73(b) Statement (when there is an assignee)  
 English Translation Document (if applicable)  
 Information Disclosure Statement (IDS) with PTO-1449  
 Preliminary Amendment  
 Return Receipt Postcard (MPEP 503)  
 Small Entity Statement(s)  
 Statement file in prior application, status still proper and desired.  
 Certified Copy of Priority Document(s) with Claim of Priority  
(if foreign priority is claimed).  
 OTHER: Check in the amount of \$760.00

If a **CONTINUING APPLICATION**, check appropriate box and supply the requisite information:

Continuation  Divisional  Continuation-in-part (CIP)  
of prior application Serial No. 08/307,588.

Amend the specification by inserting before the first line the following sentence: --This application is a  continuation,  divisional or  continuation-in-part of application Serial No. 08/307,588, filed December 5, 1994, which is in turn a National Stage of International Application PCT/EP93/00770, filed March 30, 1993--.

**CORRESPONDENCE ADDRESS:**

Foley & Lardner Address noted above.

Telephone: (202) 672-5300

Fax Number: (202) 672-5399


**FEES CALCULATIONS:** (Small entity fees indicated in parentheses.)

(1) For	(2) Number Filed	(3) Number Extra	(4) Rate	(5) Basic Fee \$760 (\$380)
Total Claims	5 - 20 =	0	x \$18 (x \$9)	
Independent Claims	2 - 3 =	0	x \$78 (x \$39)	
Assignment Recording Fee per property			\$40	
			<b>TOTAL FEE:</b>	<b>\$760.00</b>

**METHOD OF PAYMENT:**

A check in the amount of the above TOTAL FEE is attached. If payment is enclosed, this amount is believed to be correct; however, the Commissioner is hereby authorized to charge any deficiency or credit any overpayment to Deposit Account No. 19-0741.

Respectfully submitted,

  
Bernhard D. Saxe  
Reg. No. 28,665

Date: February 2, 1999  
Docket No.: 017283/0123

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

**Attorney Docket No. 017283/0123**

In re patent application of

Patrick BENOIT *et al.*

Serial No.: Unassigned

Filed: February 2, 1999

For: MONOCLONAL ANTIBODIES AGAINST THE INTERFERON RECEPTOR,  
WITH NEUTRALIZING ACTIVITY AGAINST TYPE 1 INTERFERON

**PRELIMINARY AMENDMENT**

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

Prior to examination of the above-identified application, Applicants respectfully request that the following amendments be entered:

**IN THE SPECIFICATION:**

Page 8, line 14, after "figure 3," insert --(SEQ ID NOS: 3-4)--.

Page 8, lines 18 and 25, after "figure 2," insert --(SEQ ID NOS: 1-2)--.

Page 8, line 27, after "229," insert --of SEQ ID NO: 1 or 2--.

Page 10, line 12, insert --is-- between "it" and "directed".

Page 10, line 16, after "figure 2," insert \_\_SEQ ID NOS: 1-2\_\_--.

Page 13, line 15, after "sequence" insert --(SEQ ID NOS: 1-2)--.

Page 13, line 23, after "sequence" insert --(SEQ ID NOS: 3-4)--.

Page 14, line 7, after "figure 2," insert --; SEQ ID NOS: 1-2--.

Page 24, at the end of the specification, before the claims, insert the printed Sequence Listing submitted concurrently herewith, and renumber pages 1-9 of the Sequence Listing as pages 25-33 of the specification.

**IN THE CLAIMS:**

Please delete claims 1-22 and insert the following new claims:

--23. A peptide or polypeptide which is a fragment of the extracellular portion of the IFN-R of SEQ ID NO: 2, said peptide or polypeptide consisting of amino acid residue 27 to amino acid residue 427 of SEQ ID NO: 1 or 2 or a portion thereof; wherein said peptide or polypeptide specifically binds to monoclonal antibody 64G12 (deposited at the ECACC under no. 92022605).

24. A peptide or polypeptide as claimed in claim 23, consisting of amino acid residue 27 to amino acid residue 229 of SEQ ID NO: 1 or 2 or a portion thereof.

25. A peptide or polypeptide which is a fragment of the extracellular portion of the IFN-R of SEQ ID NO: 2, said peptide or polypeptide consisting of amino acid residue 1 to amino acid residue 229 of SEQ ID NO: 1 or 2 or a portion thereof; wherein said peptide or polypeptide specifically binds to monoclonal antibody 64G12.

26. An analogue of a peptide or polypeptide as claimed in claim 23, which is derived from said peptide or polypeptide by substitution of one or more amino acid residues and which retains the ability to specifically bind to monoclonal antibody 64G12.

27. A method of producing a monoclonal antibody, comprising immunizing an animal with a peptide or polypeptide as claimed in claim 23, fusing spleen cell from the immunized animal with myeloma cells, isolating hybridoma cells which produce antibodies, and selecting and purifying monoclonal cell lines producing antibodies which specifically bind to said peptide or polypeptide.

28. A method of producing a monoclonal antibody, comprising contacting stimulated B-lymphocytes *in vitro* with a peptide or polypeptide according to claim 23, fusing the resultant B-lymphocytes with B-lymphocytes immortalized with Epstein-Barr

Attorney Docket No. 017283/0123  
Applicant: Patrick BENOIT *et al.*

virus, isolating hybridoma cells which produce antibodies, and selecting and purifying monoclonal cell lines producing antibodies which specifically bind to said peptide or polypeptide. --

**IN THE ABSTRACT**

Please insert the Abstract provided on the attached sheet.

**REMARKS**

The Examiner is respectfully requested to enter the above amendments prior to examination of the instant application. Support for the amendments is present throughout the specification, in particular at pages 10-11.

Respectfully submitted,



February 2, 1999  
Date

Bernhard D. Saxe  
Reg. No. 28,665

FOLEY & LARDNER  
3000 K Street, N.W.  
Suite 500  
Washington, D.C. 20007-5109  
Tel: (202) 672-5300

**ABSTRACT OF THE DISCLOSURE**

A monoclonal antibody is provided which is directed against the human interferon type I receptor (IFN-R), which recognizes the extracellular domain of the human IFN-R and which has neutralizing capacity against the biological properties of human type I-IFN. Diagnostic and therapeutic applications for the monoclonal antibody also are provided.

MONOCLONAL ANTIBODIES AGAINST THE INTERFERON  
RECEPTOR, WITH NEUTRALIZING ACTIVITY AGAINST  
TYPE I INTERFERON

The interferons (IFN) constitute a group of secreted proteins which exhibit a wide range of biological activities and are characterized by their capacity to induce an antiviral state in vertebrate cells (I. Gresser and M.G. Tovey Biochem Biophys. Acta 516:231, 1978). There are three antigenic classes of IFN : alpha ( $\alpha$ ), beta ( $\beta$ ) and gamma. IFN $\alpha$  and IFN $\beta$  together are known as the type I interferon.

Natural type I human interferon comprises 12 or more closely related proteins encoded by distinct genes with a high degree of structural homology (Weissmann and Weber, Prog. Nucl. Acid. Res. Mol. Biol. 33:251, 1986).

The human IFN $\alpha$  locus comprises two subfamilies. The first subfamily consists of 14 non allelic genes and 4 pseudogenes having at least 80% homology. The second subfamily,  $\alpha$ II or omega ( $\omega$ ), contains 5 pseudogenes and 1 functional gene which exhibits 70% homology with the IFN $\alpha$  genes (Weissmann and Weber 1986).

The subtypes of IFN $\alpha$  have different specific activities but they possess the same biological spectrum (Streuli et al. PNAS-USA 78:2848, 1981) and have the same cellular receptor (Agnet M. et al. in "Interferon 5" Ed. I. Gresser p. 1-22, Academic Press, London 1983).

The interferon  $\beta$  (IFN $\beta$ ) is encoded by a single gene which has approximately 50% homology with the IFN $\alpha$  genes.

The interferon  $\alpha$  subtypes and interferon  $\beta$  bind to the same receptor on the cell surface.

The interferon gamma (IFN gamma) is also encoded by a single copy, which has little homology with the IFN $\alpha$  and IFN $\beta$  genes. The receptor for IFN gamma is distinct from the receptor of the  $\alpha$  and  $\beta$  interferons.

For the purpose of the present invention the receptor of  $\alpha$  and  $\beta$  classes of IFN will be designated IFN-R. This represents natural type I receptor. The group of proteins forming natural interferon  $\alpha$  will be designated IFN $\alpha$ , and type I-IFN will represent both natural IFN $\alpha$ , IFN $\omega$ , and IFN $\beta$ .

Despite the fact that interferon is a potent antiviral agent, there is considerable evidence to suggest, that many of the characteristic symptoms of acute virus diseases such as upper respiratory tract infections are caused by an overproduction of interferon alpha. Furthermore, IFN alpha has been shown to contribute to the pathogenesis of certain chronic virus infections in experimental animals and the available evidence suggests that this is also the case for certain human chronic virus diseases such as those due to measles virus.

The interferons  $\alpha$  are also potent immuno-regulatory molecules which stimulate polyclonal B-cell activation, enhance NK cell cytotoxicity, inhibit T-cell functions, and modulate the expression of the major histocompatibility complex (MHC) class 1 antigens, all of which are implicated in the induction of autoimmunity and in graft rejection. The abnormal production of interferon  $\alpha$  is associated with a number of autoimmune diseases and inflammatory disorders including systemic lupus erythematosus (SLE), type I diabetes, psoriasis, rheumatoid arthritis, multiple sclerosis, Behçet's disease, aplastic anemia, the acquired immunodeficiency syndrome (AIDS) and severe

PCT/EP93/00770

combined immunodeficiency disease. The presence of interferon  $\alpha$  in the serum of patients with systemic lupus is correlated with both the clinical and humoral signs of increased disease activity. The production of interferon  $\alpha$  in HIV positive subjects is also highly predictive of disease evolution.

Administration of interferon  $\alpha$  has been reported to exacerbate underlying disease in patients with psoriasis and multiple sclerosis and to induce a SLE like syndrome in patients without a previous history of autoimmune disease. Interferon  $\alpha$  has also been shown to induce glomerulonephritis in normal mice and to accelerate the onset of the spontaneous autoimmune disease of NZB/W mice.

Interferon  $\alpha$  is also produced during the course of graft-versus-host disease (GVHD) in parallel with the enhanced NK cell activity characteristic of systemic GVHD. Interferon  $\alpha$  is the principal modulator of NK cell cytotoxicity and administration of interferon  $\alpha$  has been shown to enhance the intestinal consequences of GVHD in normal mice.

The object of the present invention is to provide new antagonists against the biological activities of the human type I-IFN. These antagonists could be used for therapeutical, including prophylaxis purposes, in cases where the type I-IFN (IFN  $\alpha/\beta$ ) is abnormally produced and when this abnormal production is associated with pathological symptoms. Such antagonists could also be used for the diagnosis of various diseases or for the study of the evolution of such diseases.

In order to define such antagonists, the inventors have taken into account the fact that the human natural type I-IFN is in fact constituted of a mixture of

interferons (subspecies) and the fact that the composition of this association of different subtypes of interferons varies both quantitatively and qualitatively.

Some natural interferons, such as the ones secreted by Namalwa cells (Namalwa interferon) or leukocyte (leucocyte interferon) have been studied in detail (N.B. Finter and K.H. Fautes, *Interferon 2*, 1980, p. 65-79 I. Gresser Editor Academic Press ; K. Cantell et al, *Interferon 1*, 1979 p. 2-25, I. Gresser Editor Academic Press) and were used by the inventors to define natural type I interferons.

In some pathological cases, like AIDS, interferons having some special properties have been described (O.T. Preble et al, *Annals of New-York Academy of Sciences* p. 65-75). This interferon involved in pathological cases like AIDS nevertheless binds to the same receptor, as described above.

One object of the present invention is to provide an antagonist of the type I-IFN, which would be able to inhibit or neutralize, to a determined extent, the biological properties of the human type I-IFN, that is to say, to neutralize in vivo a mixture of  $\alpha$ ,  $\beta$ ,  $\omega$  subspecies.

Accordingly the inventors have defined antibodies, especially monoclonal antibodies, which have the property of being antagonists to the type I-IFN. These antibodies are directed against the human type I-IFN receptor.

The invention thus also concerns the use of the monoclonal antibodies for the preparation of pharmaceutical compositions, useful for the treatment of symptoms associated with the abnormal production of

type I-IFN. These monoclonal antibodies are also appropriate for the preparation of diagnosis reagents.

A monoclonal antibody according to the present invention is directed against the human type I-interferon receptor (IFN-R) and is characterized by the following properties :

- it recognizes the extracellular domain of the human IFN-R, and
- it has a neutralizing capacity against the biological properties of the human type I-IFN.

The ability to neutralize the biological properties of type I-IFN can be estimated as a function of the capacity of the monoclonal antibody to neutralize the antiviral activity of the type I-IFN. Such a test is relevant in order to determine whether the antibody assayed is included within the scope of the invention, although it is clear that the biological properties of type I-IFN are not limited to its antiviral properties. Detailed procedures are given in the examples in order to enable to perform such a test of the antiviral activity. The cells tested can advantageously be Daudi-cells, which affinity for the type I-IFN is well known. The main steps of such a test would consist in :

- incubating a determined concentration of human cells responsive to human type I-IFN, with human type I-IFN in the presence of a determined concentration of monoclonal antibodies to be assayed, for a time sufficient to allow the formation of a complex between the monoclonal antibodies and the IFN-R of the human cells and/or between the type I-IFN and the IFN-R of the human cells ;

- infecting the incubated cells with a determined virus, in a determined concentration,
- washing the cells,
- resuspending the cells in culture medium,
- incubating for a time sufficient to allow virus replication ;
- lysing the cells ;
- measuring the virus replication, or measuring the inhibition of the cytopathic effect.

The ability of the monoclonal antibodies of the invention to neutralize the biological properties of the human type I-IFN can be modulated as a function of the dose of antibodies used. Accordingly a 100% inhibition of the biological properties, or a partial inhibition can be obtained.

According to another embodiment of the present invention, the monoclonal antibodies directed against the human type I-IFN receptor, are further characterized by the fact that they are capable of inhibiting the binding of a human type I-IFN, to the human IFN-R.

A monoclonal antibody having the capacity to recognize the extracellular domain of the human IFN-R and capable of inhibiting the binding of the human type I-IFN to its receptor, can be selected by the following steps :

- preincubating a determined concentration of purified monoclonal antibodies or a hybridoma culture supernatant containing monoclonal antibodies to be assayed, with human cells capable of harboring IFN-R ;
- adding labelled human type I-IFN, in a determined concentration, to the above preincubated medium ;

- incubating the medium containing the human cells, the monoclonal antibodies and the labelled type I-IFN for a time sufficient to allow an equilibrium to occur, between the monoclonal antibodies on the one hand and the type I-IFN on the other hand, with the cellular IFN-R ;
- washing the cells ;
- determining the formation of a binding complex between the human cells and the labelled type I-IFN by counting the amount of attached labelled type I-IFN.

Some of the monoclonal antibodies of the invention, have also the capacity to neutralize the antiproliferative properties of the human type I-IFN. This property can also be assayed on Daudi cells, by performing the following steps :

- allowing cells to grow in presence of human type IFN and determined concentration of mAb ;
- counting the cells in order to detect an inhibition of the antiproliferative effect of the human type I-IFN.

One property of a monoclonal antibody according to the invention resides in its capacity to recognize the extracellular domain of the human IFN receptor. This property of the monoclonal antibody can be assayed on human cells bearing the natural human receptor but also on the extracellular domain of a recombinant IFN-R such as expressed in a prokaryotic cell, for instance in E.coli or a recombinant IFN-R such as expressed in a eucaryotic cell such as mammalian cell for instance a CHO-cell.

This receptor can indeed present different properties, depending on the fact that it is produced in a prokaryotic or eucaryotic cell and accordingly

depending on the fact that the post-translational maturation occurred or not. The inventors interestingly showed that relevant assays, to evaluate the capacity of a monoclonal antibody according to the invention i.e. to recognize the cellular IFN-R, can be performed on a recombinant receptor expressed in mammalian cells. As a matter of fact, such recombinant receptor has the same properties as the cellular receptor, as far as its recognizing activity is concerned.

Monoclonal antibodies of the invention can be obtained against various forms of the receptor, including the complete receptor, a particular domain or a peptide characteristic of the aminoacid sequence of the receptor represented in figure 3.

Monoclonal antibodies of the invention can for example be prepared against the soluble form of the receptor. A hydrosoluble polypeptide corresponding to the soluble form of the INF-R is described on figure 2. According to the present invention, a soluble form of the IFN-R corresponds to a peptide or a polypeptide, capable of circulating in the body.

Other monoclonal antibodies according to the invention can also be prepared against a peptide comprised in the extracellular domain of the receptor as described on figure 2. An advantageous peptide corresponds for instance to the aminoacid sequence comprised between aminoacid 1 and aminoacid 229. According to another embodiment of the invention, the antibodies can be prepared against a polypeptide modified by substitution of one or more amino acids, provided that antibodies directed against the non modified extracellular domain of the IFN-R, recognize the modified polypeptide or peptide.

Preferred monoclonal antibodies according to the invention are those which are of the IgG1 type.

Among the antibodies of the invention, an antibody which has the capacity of inhibiting the binding of the type I-IFN to its receptor is preferably characterized in that it inhibits the in vitro binding of human type IFN, to the human cellular IFN-R when it is co-incubated with cells harboring the hu-IFN-R, at a concentration of antibodies equal or inferior to 100  $\mu$ g/ml, preferably equal or inferior to 50  $\mu$ g/ml, advantageously inferior to 20  $\mu$ g/ml, more preferably in the range of approximately 0.5 to 2  $\mu$ g/ml.

The inventors have shown that the high affinity binding capacity of a monoclonal antibody is not sufficient to ensure that this antibody will be able to inhibit the binding activity of the human type I-IFN to the IFN-R. Nevertheless the high affinity binding capacity of the monoclonal antibody is necessary to investigate further the ability of the antibody to inhibit the binding of the type I-IFN to its cellular receptor.

Another monoclonal antibody is characterized in that it neutralizes in vitro the antiproliferative activity of human type I-IFN, on cells highly responsive to this human type I-IFN, for instance Daudi cells at a concentration in a range of 1 to 10  $\mu$ g/ml.

According to another embodiment a monoclonal antibody is also characterized in that it neutralizes in vitro the antiproliferative activity of human type IFN, on cells poorly responsive to this human IFN, for instance Ly28 cells, at a concentration in a range of 50 to 100  $\mu$ g/ml.

A particular group of monoclonal antibodies according to the invention is characterized in that it

neutralizes the antiviral activity of the human type I-IFN, on cells highly responsive to this human type I-IFN, for instance Daudi cells at a concentration in a range of 1 to 50  $\mu$ g/ml, preferably 1 to 20  $\mu$ g/ml, for a concentration of type I-IFN in the range of 1 to 1000 units with reference to the international standard MRC 69/19.

Advantageously, the monoclonal antibody according to the invention is such that these antibodies do not bind to the human receptor for IFN gamma.

One particular antibody satisfying the requirements of the invention, is such as it directed against an epitope on the amino-acid sequence comprised between amino-acid 27 and amino-acid 427 of the extracellular domain of the human IFN-R as represented on figure 2.

One particularly interesting monoclonal antibody according to the invention is the antibody designated 64G12 under n° 92022605 which has been deposited at the ECACC (European Collection of Animal Cell Cultures Porton Down Salisbury, Wiltshire SP4 056, United Kingdom) on February 26, 1992.

These antibodies may be prepared by conventional methods involving the preparation of hybridoma cells by the fusion of myeloma cells and spleen cells of an animal immunized beforehand with the peptide antigen, on the conditions such that the antigen against which the antibodies are formed is constituted by the extracellular domain of IFN-R or any polypeptide or peptide of this domain.

The hybridomas are constructed according to the protocole of Kohler and Milstein (Nature, 1974, 256: 495-497). For example the hybridomas are derived from

the fusion of the spleen cells above described with NS1 mouse (BalbC) HGPRT<sup>+</sup> as myeloma cell.

A second procedure for the production of monoclonal antibodies according to the invention, consists in carrying out the fusion between B-cells of blood immortalized with the Epstein/Barr virus and human B lymphocytes placed beforehand in contact with the extracellular domain or a fragment thereof of the IFN-R, against which it is decided to form monoclonal antibodies. B-cells placed in contact beforehand with the extracellular domain of IFN-R or fragment thereof against which it is decided to form monoclonal antibodies, may be obtained by in vitro culture contacted with the antigens, the recovery of the B-cells coated with these antigens being preceded by one or several cycles of stimulation.

The invention thus concerns human antibodies as obtained by carrying out the above procedure, having the above defined properties.

The invention also aims at providing a monoclonal antibody characterized in that the variable or complementary determining regions of its heavy and/or light chains are grafted on the framework and/or constant regions of a human antibody.

The invention further provides a composition having antagonist properties for the biological properties of the human type I-IFN, characterized in that it comprises monoclonal antibodies as defined above.

Accordingly the invention provides a pharmaceutical composition characterized in that it comprises monoclonal antibodies as defined above, together with an appropriate pharmaceutical vehicle.

The invention also concerns the use of a monoclonal antibody as defined above, for the manufacture of a drug for the treatment or profilaxis of a pathological state or symptoms associated with overproduction of type-I-IFN.

According to a first example, the antibodies can be used in a pharmaceutical composition, for the treatment of allograft rejection.

According to another example, antibodies of the invention are used as active principle in a pharmaceutical composition for the treatment of autoimmune and inflammatory diseases. Such diseases include systemic lupus erythematosus, type 1 diabetes, psoriasis, rheumatoid arthritis, multiple sclerosis, Behçet's disease, aplastic anemia, acquired immunodeficiency syndrome (AIDS), and severe combined immunodeficiency disease.

Treatment of acute virus diseases can also be performed with the antibodies of the invention. As example upper respiratory tract infections, chronic virus infections such as those due to measles virus, can be performed.

The antibodies of the invention can also be used for the in vitro diagnosis of the presence of the human type I-IFN receptor or cells.

Further details and additional information will arise from the description from the description of the examples and from the figures.

FIGURES

- Figure 1 : binding of  $^{125}\text{I}$ -labelled monoclonal antibodies 34F10 and 64G12 to :

- A : Daudi cells
- B : Ly28 cells

Briefly,  $10^6$  cells were incubated for 2 hours at 4°C in presence of different amounts of the labelled antibodies diluted in RPMI medium containing 10% fetal calf serum (FCS). The cells were then washed 4 times in RPMI-1% FCS and counted for bound radioactivity. Nonspecific binding was measured by incubation with a 100 fold excess of cold antibodies and subtracted from total counts.

- Figure 2 : nucleotide and corresponding amino-acid sequence of the extracellular domain of the human IFN-R

The monoclonal antibodies were produced against recombinant soluble forms of the human interferon alpha-beta receptor (IFN-R) synthesized in either prokaryotic cells (E.coli) or a mammalian cell system (Cos cell). These soluble forms were based on the DNA sequence described in figure 2.

- Figure 3 : nucleotide and corresponding amino-acid sequence of the human IFN-R.

## EXAMPLES

**EXAMPLE 1 :**

## Synthesis of the soluble receptors

### Synthesis in *E.coli*

A fragment of DNA containing the sequence coding for the extracellular domain (amino acids 27 to 427) of the human INF-R (figure 2), in which an extra-sequence coding for 5 histidyl residues was introduced just before the termination codon, was cloned in the expression vectors pKK233-2. This fragment was produced by the Polymerase Chain Reaction (PCR) and the resulting plasmids were sequenced to confirm both in-frame insertion with the Shine-Dalgarno sequence and the appropriate sequence coding for the receptor.

The poly-histidyl tail introduced into the recombinant protein enables it to be purified rapidly by affinity chromatography on a chelated nickel support (NTA column) as described previously (Hochuli E. et al, Bio/technology, 1988, 1321-1325).

The plasmid was introduced into the E.coli strain, JM105, and protein synthesis induced by addition of IPTG to the culture medium (pKK233-2, tac promoter).

Proteins were extracted from the bacterial pellet and the soluble receptor purified to homogeneity by affinity chromatography as described hereafter. This procedure yielded a protein that migrates as 2 bands around 50 kDa under reducing conditions and three bands under non-reducing conditions. The maximum concentration of the protein obtained by different procedures was approximately 20 $\mu$ g/ml.

The N-terminal sequence of the two proteins detected by gel electrophoresis has shown that both proteins are the expected fragment of the receptor.

Synthesis and purification of an unglycosylated soluble receptor :

Bacterial culture (250ml)

|  
|

IPTG induction 3h

|  
|

cell pellet

6M Guanidine hydrochloride pH8

|  
|

centrifugation

|  
|

NTA column:

Washes pH 8 urea 8M

|  
|

pH 6,3 urea 8M

pH 5,9 urea 8M

Elution pH 4 urea 8M

|  
|

refolding dilution, dialysis

|  
|

against Tris 0,1 M pH9

dialysis PBS

Using the same PCR approach, we also constructed an expression vector coding for the IFN-R amino acid sequence 1-427, with an additional 5-histidyl residues at the C-terminus, inserted in expression vector pXMT-3. The exact nucleotide sequence of the insert was also confirmed.

The resulting plasmid was introduced by electroporation into Cos7 cells for transient expression and the recombinant protein was purified to homogeneity by affinity chromatography followed by ion exchange chromatography on mono-Q (Pharmacia) as described hereafter.

## Purification of the soluble IFN- $\beta$ from Cos7 cells

## preparative electroporation of cos cells

18 h

serum free medium

1

supernatants taken after 48h, 72h, 96h

1

## concentration

1

### NTA column

100

Wash PBS

elution 0.1 M NaOAc pH 5.5

— 1 —

### neutralization

concentration, 30 000 cut off

11

Mono Q (0-0.5 M Na Cl)

This purification yielded to a 76 kDa protein whose N-terminal sequence corresponds to the predicted receptor sequence with some heterogeneity in the processing of the leader sequence.

**EXAMPLE 2 :**

Production of monoclonal antibodies against the interferon type I receptor

1) Production of the monoclonal antibodies

Mice were immunized by injection of recombinant soluble interferon (r sIFN-R) purified from E.coli or from a culture supernatant of Cos7 cells. Initially mice were injected both intraperitoneally and subcutaneously with the purified protein in complete Freund's adjuvant. Subsequently mice were injected once a week intraperitoneally with the purified proteins diluted in buffered saline solution. Ten micrograms of recombinant proteins were injected each time.

After the fourth injection, blood was collected and the presence of specific serum antibodies were tested by both ELISA and Western blot against the recombinant receptor. The strongest responders were then boosted with a total of 10 $\mu$ g of antigen half of which was injected intravenously and half intraperitoneally.

PCT/EP93/00770

2) Cell fusion

Four days after boosting, spleen cells from the immunized animal were collected and fused to NS1 (mouse) (Balbc) HGPRT<sup>+</sup> myeloma cells according to the method described by S. Fazekas et al. (J. Immunol. Methods 35:1-32, 1980). Briefly,  $5 \times 10^7$  spleen cells were fused to  $3 \times 10^7$  myeloma cells in 1ml of polyethylene glycol solution and distributed in five 96 well plates on a peritoneal macrophage feeder layer in HAT (hypoxanthine, aminoprotein and thymidine) medium. This procedure was repeated 4 times as  $20 \times 10^7$  spleen cells were obtained from the immunized mouse. Screening for specific hybridomas was undertaken when large colonies were detectable in culture wells.

For the screening, presence of specific antibodies was determined by a direct ELISA method :

a) ELISA plates were coated overnight at 4°C with purified E.coli-expressed or Cos7 cell-expressed sIFN- $\beta$  diluted in PBS. Plates coated with BSA were used to detect non specific binding,

b) Plates were saturated by incubation with 3% BSA in PBS for 1 hour at 37°C,

c) Plates were incubated for 4 hours at room temperature with hybridoma supernatants diluted 1 in 4 with PBS-0.05% Tween 20,

d) Bound antibodies were detected by a two step procedure, comprising a first incubation with goat anti-mouse biotinylated immunoglobulin followed by streptavidin-horseradish peroxidase complex (both from Amersham and diluted 1/1000 in PBS-0.05% Tween 20).

Positive antibody secreting hybridomas were passaged in 24 well plates on a spleen cell feeder layer and their reactivity was again checked by ELISA, and Western-blot.

3) Identification of reactivity to the natural interferon type I receptor

The reactivity of the monoclonal antibodies (mAbs) recognizing the recombinant sIFN-R was tested against the natural class I receptor expressed at the surface of Daudi cells, by membrane immunofluorescence. Briefly,  $5 \times 10^5$  Daudi cells were incubated in 100 $\mu$ l of culture supernatant of chosen hybridomas for 30 min at 4°C. The cells were then washed 4 times in RPMI medium containing 1% BSA and further incubated with a diluted FITC labelled goat anti-mouse F(ab')<sub>2</sub> for 30 min at 4°C. The cells were finally analyzed by flow cytometry after washing. One of the 35 tested antibodies produced against the E.coli recombinant receptor and 5 of the 6 tested antibodies produced against the COS recombinant receptor were found to recognize the natural receptor on the Daudi cells.

Cloning of these hybridomas was then performed by limiting dilution. The isotype of these mAbs was determined by an ELISA method using isotype specific antibodies. All 6 mAbs were found to be IgG1 with kappa light chains. A summary of the reactivity of these 6 mAbs is given in Table 1.

Monoclonal antibodies were purified from culture supernatants by protein G chromatography.

Table 1 :Reactivity of the anti IFN-R monoclonal antibodies

	Reactivity against the recombinant receptor				Reactivity against * the cellular receptor	
	E.COLI		COS			
	ELISA	Western	ELISA	Western		
34F10	+	+	+	+	+	
64G12	+	+	+	+	+	
63F6 64G2 64D10 65D8	-	-	+	+	weak	

\* measured on Daudi cells

EXAMPLE 3 :Inhibition of the binding of interferon to human cell lines

Inhibition of interferon binding to human cells was assayed as follows.  $10^6$  cells were preincubated at 4°C for 30 min with various dilutions of hybridoma culture supernatants or purified mAbs or with medium alone.  $^{125}$ I-labelled IFN alpha 8 or alpha 2 was added at the concentration of 100pM and cells incubated for a further 2 hours at 4°C. These incubations were performed in RPMI medium containing 20mM HEPES pH 7.4 and 10% foetal calf serum (FCS). The cells were finally washed 4 times with RPMI - 1% FCS and counted to determine bound radioactivity.

The mAb secreted by the hybridoma line 64G12 (latter named mAb 64G12) was shown in this assay to inhibit the binding of labelled IFN to the cells in a dose-dependent manner. 50% inhibition of binding to the Daudi cells (Burkitt lymphoma cell line ; Klein et al., Cancer Researh, 28:1300-1310, 1968) was obtained at a mAb concentration of 0.4 $\mu$ g/ml. The same inhibition was obtained with K562 cells (chronic myelogenous leukemia, Lozzio and Lozzio, Cell, 45:321-334, 1975) but 50% inhibition was obtained at 11 $\mu$ g/ml for HL60 cells (Promyelocytic leukemia, Collins S.J. et al., Nature, 270:347-349, 1977) and 60 $\mu$ g/ml for Ly28 cells (Klein G. et al. Int. J. Cancer, 10:44-57, 1972).

Table 2 :

The inhibition of binding of labelled IFN alpha 2 to various cell lines by mAB64G12

Cell liner	Concentration of mAB which gives 50% inhibition of binding
Daudi K562	0,4 $\mu$ g/ml
HL60	11 $\mu$ g/ml
Ly28	60 $\mu$ g/ml

The difference in the mAb concentration at which 50% inhibition of binding of IFN is obtained has been investigated by direct binding of  $^{125}$ I-labelled mABs 64G12 and 34F10 to the same cell lines and Scatchard

plot analysis of the results. In the concentration range of 0.1 to 1.5  $\mu$ g/ml, a high affinity binding of the mAb 34F10 ( $\approx$ 10nM) was seen on all cell lines whereas a high affinity binding of mAb 64G12 was only detected on Daudi and K562 cells (Figure 1).

EXAMPLE 4 :

Inhibition of the function of type I interferon

Functional inhibition of type I interferon by the purified mAb 64G12 was demonstrated in an antiviral assay on Daudi cells using either recombinant IFN alpha 2, IFN beta and IFN omega, or purified Namalwa and leucocyte interferons, and in an antiproliferative assay with recombinant IFN alpha 2.

\* Antiviral activity

An antiviral assay on Daudi cells was performed as described (M. Dron and M.G. Tovey, J. Gen. Virol. 64:2641-2647, 1983). Cells ( $0.5 \times 10^6$ /ml) were incubated for 24 hours in the presence of interferon and antibodies.  $10^6$  cells in 1 ml were then infected for 1 hour at 37°C with Vesicular stomatitis virus (VSV) then washed 3 times, resuspended in culture medium and incubated for 18 hours at 37°C. Cells were then lysed by freeze-thawing and virus replication measured by titration of the supernatants on L929 cells. A dose-dependent inhibition of the antiviral activity of the various subtypes of type I IFN was demonstrated for the purified mAb 64G12.

For the antiviral assay with the Wish cells, cells were incubated for 24 hours with various concentrations of interferons in the presence of the mAbs prior to challenge with VSV. In this assay, the mAb 64G12 was demonstrated to block completely the antiviral activity of Leukocyte IFN (50U/ml), recombinant IFN alpha 2 (50U/ml) and interferon from the sera of AIDS patients (50, 75 and 150U/ml).

**\* antiproliferative activity**

For the antiproliferative assay, Daudi cells were seeded at a concentration of  $10^5$  cells per ml in a 96 well plate in the presence of interferon and purified inhibitory or control antibody. Cells were then counted after 24, 48 and 72 hours with a Coulter counter and checked for viability by trypan blue exclusion. Purified mAb 64G12 demonstrated a dose-dependent inhibition of the antiproliferative activity of interferon alpha 2.

CLAIMS

1. Monoclonal antibody directed against the human interferon class I receptor (IFN-R) characterized by the following properties :
  - it recognizes the extracellular domain of the human IFN-R, and
  - it has a neutralizing capacity against the biological properties of the human type I-IFN.
2. Monoclonal antibody directed against the human type I IFN-R according to claim 1, characterized by its capacity to inhibit the binding of a human pathological type I-IFN, to the IFN-R.
3. Monoclonal antibody according to claim 1 or 2, which is obtainable from a hybridoma cell prepared by fusion of a myeloma cell with spleen cells from an animal previously immunized with the soluble form of the human IFN-R.
4. Monoclonal antibody according to anyone of claims 1, 2 or 3, characterized in that it recognizes an epitope on a soluble form of the human cellular IFN-R or of a recombinant IFN-R.
5. Monoclonal antibody according to anyone of claims 1 to 4, characterized in that it inhibits in vitro the binding of human type I-IFN, to the human cellular IFN-R when it is co-incubated with cells harboring the hu-IFN-R, at a concentration of antibodies equal or inferior to 100 µg/ml, preferably equal or inferior to 50 µg/ml, advantageously inferior to 20 µg/ml, more preferably in the range of approximately 0,5 to 2 µg/ml.
6. Monoclonal antibody according to anyone of claims 1 to 5, characterized in that it neutralizes in vitro the antiproliferative activity of the human type I-IFN, on cells highly responsive to this human type I-IFN,

for instance Daudi cells at a concentration in a range of 1 to 10  $\mu$ g/ml.

7. Monoclonal antibody according to anyone of claims 1 to 6, characterized in that it neutralizes in vitro the antiproliferative activity of human type I-IFN, on cells poorly responsive to this human type I-IFN, for instance Ly28 cells, at a concentration in a range of 50 to 100  $\mu$ g/ml.

8. Monoclonal antibody according to anyone of claims 1 to 7, characterized in that it does not bind to the human receptor of the IFN gamma.

9. Monoclonal antibody according to anyone of claims 1 to 8, characterized in that it recognizes an epitope on the aminoacid sequence 27 to 427 of the human IFN-R.

10. Monoclonal antibody according to anyone of claims 1 to 9, characterized in that it neutralizes in vitro the antiviral activity of the human type I-IFN, on cells highly responsive to this human type I-IFN, for instance Daudi cells at a concentration in a range of 1 to 10  $\mu$ g/ml.

11. Monoclonal antibody according to anyone of claims 1 to 10, characterized in that it neutralizes in vitro the antiviral activity of the human class I-IFN, on cells poorly responsive to this human IFN, for instance Ly28 cells, at a concentration in a range of 50 to 100  $\mu$ g/ml.

12. Monoclonal antibody according to anyone of claims 1 to 11, characterized in that it is the 64G12 antibody, deposited at the ECACC on February 26, 1992 under n° 92022605.

13. Monoclonal antibody according to anyone of claims 1 to 11, characterized in that it is a humanized antibody, for instance characterized in that the variable or complementary determining regions of its

PCT/EP93/00770

heavy and light chains are grafted on the framework and constant regions of a human antibody.

14. Monoclonal antibody according to anyone of claims 1 to 11, characterized in that it is a human antibody.
15. Monoclonal antibody according to anyone of claims 1 to 11, characterized in that it is an IgG1 type antibody.
16. Hybridoma cell, characterized in that it produces monoclonal antibodies according to claims 1 to 13.
17. Composition having antagonist properties to the type I-IFN, characterized in that it comprises monoclonal antibodies according to anyone of claims 1 to 16.
18. Pharmaceutical composition, characterized in that it comprises monoclonal antibodies according to anyone of claims 1 to 17, together with an appropriate pharmaceutical vehicle.
19. Use of a monoclonal antibody according to anyone of claims 1 to 17, for the manufacture of a drug for the treatment or prophylaxis of a pathological state associated with proliferative cell activity and/or viral cell infection.
20. Process for the selection of a monoclonal antibody having the capacity to recognize the extracellular domain of the human IFN-R and capable of inhibiting the binding of the human type I-IFN, to the IFN-R, characterized by the following steps :
  - preincubating a determined concentration of purified monoclonal antibodies according to anyone of claims 1 to 15 or a hybridoma culture supernatant containing monoclonal antibodies, with human cells susceptible of harboring IFN-R ;
  - adding labelled human type I-IFN in a determined concentration, to the above preincubating medium ;

卷之三

- incubating the medium containing the human cells, monoclonal antibodies and labelled type I-IFN for a time sufficient to allow an equilibrium to occur, between the monoclonal antibodies on the one hand and the type I-IFN on the other hand, with the cellular IFN-R ;
- washing the cells ;
- determining the formation of a binding complex between the human cells and the type I-IFN, by counting the amount of attached labelled type I-IFN.

21. Process for the selection of a monoclonal antibody having the capacity to recognize the extra-cellular domain of the human IFN-R and having a neutralizing capacity against the antiproliferative activities of the type I-IFN, on human cells characterized by the steps of :

- allowing cells to grow in the presence of human type I-IFN and in the presence of a determined concentration of monoclonal antibodies according to anyone of claims 1 to 15 ;
- counting the cells in order to detect an inhibition of the antiproliferative effect of the type I-IFN.

22. Process for the selection of a monoclonal antibody having the capacity to recognize the extra-cellular domain of the human IFN-R and having a neutralizing capacity against the antiviral activities of the natural, non pathological or pathological type I-IFN on human cells, characterized by the steps of :

- incubating cells with type I-IFN and monoclonal antibodies according to anyone of claims 1 to 15, in determined concentrations, for a time sufficient to allow the formation of a complex

between the monoclonal antibodies and the IFN-R of the human cells and/or between the type I-IFN and the IFN-R of the human cells ;

- infecting the above incubated cells with a determined concentration of a virus ;
- washing the cells ;
- resuspending the cells in culture medium ;
- incubating for a time sufficient to allow the replication of the virus ;
- lysing the cells and ;
- measuring the virus replication or measuring the inhibition of the cytopathic effect.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: BENOIT, Patrick  
MEYER, Francois  
MAGUIRE, Deborah  
PLAVEC, Ivan  
TOVEY, Michael G.
- (ii) TITLE OF INVENTION: MONOCLONAL ANTIBODIES AGAINST THE INTERFERON RECEPTOR, WITH NEUTRALIZING ACTIVITY AGAINST TYPE I INTERFERON
- (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Foley & Lardner
  - (B) STREET: 3000 K Street, N.W., Suite 500
  - (C) CITY: Washington
  - (D) STATE: D.C.
  - (E) ZIP: 20007
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 08/307,588
  - (B) FILING DATE: 05-DEC-1994
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: PCT/EP93/00770
  - (B) FILING DATE: 30-MAR-1993
- (viii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: EP 92400902.0
  - (B) FILING DATE: 31-MAR-1992
- (ix) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: SAXE, Bernhard D.
  - (B) REGISTRATION NUMBER: 28,665
  - (C) REFERENCE/DOCKET NUMBER: 17283/117/GUPL
- (x) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (202) 672-5300
  - (B) TELEFAX: (202) 672-5399

## (2) INFORMATION FOR SEQ ID NO: 1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1343 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 27..1334

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CTGCAGGGAT	CTGCGGCCGC	TCCCCAG	ATG ATG GTC GTC CTC CTG GGC GCG ACG	53											
Met	Met	Val	Val	Leu	Leu	Gly	Ala	Thr							
1						5									
ACC CTA GTG CTC GTC GCC GTG GGC CCA TGG GTG TTG TCC GCA GCC GCA	101														
Thr	Leu	Val	Leu	Val	Ala	Val	Gly	Pro	Trp	Val	Leu	Ser	Ala	Ala	Ala
10		15				20					25				
GGT GGA AAA AAT CTA AAA TCT CCT CAA AAA GTA GAG GTC GAC ATC ATA	149														
Gly	Gly	Lys	Asn	Leu	Lys	Ser	Pro	Gln	Lys	Val	Glu	Val	Asp	Ile	Ile
30				35					40						
GAT GAC AAC TTT ATC CTG AGG TGG AAC AGG AGC GAT GAG TCT GTC GGG	197														
Asp	Asp	Asn	Phe	Ile	Leu	Arg	Trp	Asn	Arg	Ser	Asp	Glu	Ser	Val	Gly
45				50					55						
AAT GTG ACT TTT TCA TTC GAT TAT CAA AAA ACT GGG ATG GAT AAT TGG	245														
Asn	Val	Thr	Phe	Ser	Phe	Asp	Tyr	Gln	Lys	Thr	Gly	Met	Asp	Asn	Trp
60				65					70						
ATA AAA TTG TCT GGG TGT CAG AAT ATT ACT AGT ACC AAA TGC AAC TTT	293														
Ile	Lys	Leu	Ser	Gly	Cys	Gln	Asn	Ile	Thr	Ser	Thr	Lys	Cys	Asn	Phe
75				80					85						
TCT TCA CTC AAG CTG AAT GTT TAT GAA GAA ATT AAA TTG CGT ATA AGA	341														
Ser	Ser	Leu	Lys	Leu	Asn	Val	Tyr	Glu	Glu	Ile	Lys	Leu	Arg	Ile	Arg
90				95					100				105		
GCA GAA AAA GAA AAC ACT TCT TCA TGG TAT GAG GTT GAC TCA TTT ACA	389														
Ala	Glu	Lys	Glu	Asn	Thr	Ser	Ser	Trp	Tyr	Glu	Val	Asp	Ser	Phe	Thr
110					115					120					
CCA TTT CGC AAA GCT CAG ATT GGT CCT CCA GAA GTA CAT TTA GAA GCT	437														
Pro	Phe	Arg	Lys	Ala	Gln	Ile	Gly	Pro	Pro	Glu	Val	His	Leu	Glu	Ala
125					130					135					
GAA GAT AAG GCA ATA GTG ATA CAC ATC TCT CCT GGA ACA AAA GAT AGT	485														
Glu	Asp	Lys	Ala	Ile	Val	Ile	His	Ile	Ser	Pro	Gly	Thr	Lys	Asp	Ser
140					145					150					
GTT ATG TGG GCT TTG GAT GGT TTA AGC TTT ACA TAT AGC TTA CTT ATC	533														
Val	Met	Trp	Ala	Leu	Asp	Gly	Leu	Ser	Phe	Thr	Tyr	Ser	Leu	Leu	Ile
155					160					165					
TGG AAA AAC TCT TCA GGT GTA GAA GAA AGG ATT GAA AAT ATT TAT TCC	581														
Trp	Lys	Asn	Ser	Ser	Gly	Val	Glu	Glu	Ile	Glu	Asn	Ile	Tyr	Ser	
170					175					180			185		
AGA CAT AAA ATT TAT AAA CTC TCA CCA GAG ACT ACT TAT TGT CTA AAA	629														
Arg	His	Lys	Ile	Tyr	Lys	Leu	Ser	Pro	Glu	Thr	Thr	Tyr	Cys	Leu	Lys
190					195					200					
GTT AAA GCA GCA CTA CTT ACG TCA TGG AAA ATT GGT GTC TAT AGT CCA	677														
Val	Lys	Ala	Ala	Leu	Leu	Thr	Ser	Trp	Lys	Ile	Gly	Val	Tyr	Ser	Pro
205					210					215					
GTA CAT TGT ATA AAG ACC ACA GTT GAA AAT GAA CTA CCT CCA CCA GAA	725														
Val	His	Cys	Ile	Lys	Thr	Thr	Val	Glu	Asn	Glu	Leu	Pro	Pro	Pro	Glu
220					225					230					
AAT ATA GAA GTC AGT GTC CAA AAT CAG AAC TAT GTT CTT AAA TGG GAT	773														
Asn	Ile	Glu	Val	Ser	Val	Gln	Asn	Gln	Tyr	Val	Leu	Lys	Trp	Asp	
235					240					245					

TAT ACA TAT GCA AAC ATG ACC TTT CAA GTT CAG TGG CTC CAC GCC TTT Tyr Thr Tyr Ala Asn Met Thr Phe Gln Val Gln Trp Leu His Ala Phe 250 255 260 265	821
TTA AAA AGG AAT CCT GGA AAC CAT TTG TAT AAA TGG AAA CAA ATA CCT Leu Lys Arg Asn Pro Gly Asn His Leu Tyr Lys Trp Lys Gln Ile Pro 270 275 280	869
GAC TGT GAA AAT GTC AAA ACT ACC CAG TGT GTC TTT CCT CAA AAC GTT Asp Cys Glu Asn Val Lys Thr Thr Gln Cys Val Phe Pro Gln Asn Val 285 290 295	917
TTC CAA AAA GGA ATT TAC CTT CTC CGC GTA CAA GCA TCT GAT GGA AAT Phe Gln Lys Gly Ile Tyr Leu Leu Arg Val Gln Ala Ser Asp Gly Asn 300 305 310	965
AAC ACA TCT TTT TGG TCT GAA GAG ATA AAG TTT GAT ACT GAA ATA CAA Asn Thr Ser Phe Trp Ser Glu Glu Ile Lys Phe Asp Thr Glu Ile Gln 315 320 325	1013
GCT TTC CTA CTT CCT CCA GTC TTT AAC ATT AGA TCC CTT AGT GAT TCA Ala Phe Leu Leu Pro Pro Val Phe Asn Ile Arg Ser Leu Ser Asp Ser 330 335 340 345	1061
TTC CAT ATC TAT ATC GGT GCT CCA AAA CAG TCT GGA AAC ACG CCT GTG Phe His Ile Tyr Ile Gly Ala Pro Lys Gln Ser Gly Asn Thr Pro Val 350 355 360	1109
ATC CAG GAT TAT CCA CTG ATT TAT GAA ATT ATT TTT TGG GAA AAC ACT Ile Gln Asp Tyr Pro Leu Ile Tyr Glu Ile Ile Phe Trp Glu Asn Thr 365 370 375	1157
TCA AAT GCT GAG AGA AAA ATT ATC GAG AAA AAA ACT GAT GTT ACA GTT Ser Asn Ala Glu Arg Lys Ile Ile Glu Lys Lys Thr Asp Val Thr Val 380 385 390	1205
CCT AAT TTG AAA CCA CTG ACT GTA TAT TGT GTG AAA GCC AGA GCA CAC Pro Asn Leu Lys Pro Leu Thr Val Tyr Cys Val Lys Ala Arg Ala His 395 400 405	1253
ACC ATG GAT GAA AAG CTG AAT AAA AGC AGT GTT TTT AGT GAC GCT GTA Thr Met Asp Glu Lys Leu Asn Lys Ser Ser Val Phe Ser Asp Ala Val 410 415 420 425	1301
TGT GAG AAA ACA AAA CCA GGA AAT ACC TCT AAA TGAGGTACC Cys Glu Lys Thr Lys Pro Gly Asn Thr Ser Lys 430 435	1343

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 436 amino acids
- (B) TYPE: amino acid
- (C) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Met Val Val Leu Leu Gly Ala Thr Thr Leu Val Leu Val Ala Val 1 5 10 15
Gly Pro Trp Val Leu Ser Ala Ala Gly Gly Lys Asn Leu Lys Ser 20 25 30

Pro Gln Lys Val Glu Val Asp Ile Ile Asp Asp Asn Phe Ile Leu Arg  
35 40 45

Trp Asn Arg Ser Asp Glu Ser Val Gly Asn Val Thr Phe Ser Phe Asp  
50 55 60

Tyr Gln Lys Thr Gly Met Asp Asn Trp Ile Lys Leu Ser Gly Cys Gln  
65 70 75 80

Asn Ile Thr Ser Thr Lys Cys Asn Phe Ser Ser Leu Lys Leu Asn Val  
85 90 95

Tyr Glu Glu Ile Lys Leu Arg Ile Arg Ala Glu Lys Glu Asn Thr Ser  
100 105 110

Ser Trp Tyr Glu Val Asp Ser Phe Thr Pro Phe Arg Lys Ala Gln Ile  
115 120 125

Gly Pro Pro Glu Val His Leu Glu Ala Glu Asp Lys Ala Ile Val Ile  
130 135 140

His Ile Ser Pro Gly Thr Lys Asp Ser Val Met Trp Ala Leu Asp Gly  
145 150 155 160

Leu Ser Phe Thr Tyr Ser Leu Leu Ile Trp Lys Asn Ser Ser Gly Val  
165 170 175

Glu Glu Arg Ile Glu Asn Ile Tyr Ser Arg His Lys Ile Tyr Lys Leu  
180 185 190

Ser Pro Glu Thr Thr Tyr Cys Leu Lys Val Lys Ala Ala Leu Leu Thr  
195 200 205

Ser Trp Lys Ile Gly Val Tyr Ser Pro Val His Cys Ile Lys Thr Thr  
210 215 220

Val Glu Asn Glu Leu Pro Pro Pro Glu Asn Ile Glu Val Ser Val Gln  
225 230 235 240

Asn Gln Asn Tyr Val Leu Lys Trp Asp Tyr Thr Tyr Ala Asn Met Thr  
245 250 255

Phe Gln Val Gln Trp Leu His Ala Phe Leu Lys Arg Asn Pro Gly Asn  
260 265 270

His Leu Tyr Lys Trp Lys Gln Ile Pro Asp Cys Glu Asn Val Lys Thr  
275 280 285

Thr Gln Cys Val Phe Pro Gln Asn Val Phe Gln Lys Gly Ile Tyr Leu  
290 295 300

Leu Arg Val Gln Ala Ser Asp Gly Asn Asn Thr Ser Phe Trp Ser Glu  
305 310 315 320

Glu Ile Lys Phe Asp Thr Glu Ile Gln Ala Phe Leu Leu Pro Pro Val  
325 330 335

Phe Asn Ile Arg Ser Leu Ser Asp Ser Phe His Ile Tyr Ile Gly Ala  
340 345 350

Pro Lys Gln Ser Gly Asn Thr Pro Val Ile Gln Asp Tyr Pro Leu Ile  
355 360 365

Tyr Glu Ile Ile Phe Trp Glu Asn Thr Ser Asn Ala Glu Arg Lys Ile  
370 375 380

Ile Glu Lys Lys Thr Asp Val Thr Val Pro Asn Leu Lys Pro Leu Thr  
385 390 395 400  
Val Tyr Cys Val Lys Ala Arg Ala His Thr Met Asp Glu Lys Leu Asn  
405 410 415  
Lys Ser Ser Val Phe Ser Asp Ala Val Cys Glu Lys Thr Lys Pro Gly  
420 425 430  
Asn Thr Ser Lys  
435

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 1755 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 27..1697

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CTGCAGGGAT CTGCGGCGGC TCCCCAG ATG ATG GTC GTC CTC CTG GGC GCG ACG 53  
Met Met Val Val Leu Leu Gly Ala Thr  
1 5  
ACC CTA GTG CTC GTC GCC GTG GGC CCA TGG GTG TTG TCC GCA GCC GCA 101  
Thr Leu Val Leu Val Ala Val Gly Pro Trp Val Leu Ser Ala Ala Ala  
10 15 20 25  
GGT GGA AAA AAT CTA AAA TCT CCT CAA AAA GTA GAG GTC GAC ATC ATA 149  
Gly Gly Lys Asn Leu Lys Ser Pro Gln Lys Val Glu Val Asp Ile Ile  
30 35 40  
GAT GAC AAC TTT ATC CTG AGG TGG AAC AGG AGC GAT GAG TCT GTC GGG 197  
Asp Asp Asn Phe Ile Leu Arg Trp Asn Arg Ser Asp Glu Ser Val Gly  
45 50 55  
AAT GTG ACT TTT TCA TTC GAT TAT CAA AAA ACT GGG ATG GAT AAT TGG 245  
Asn Val Thr Phe Ser Phe Asp Tyr Gln Lys Thr Gly Met Asp Asn Trp  
60 65 70  
ATA AAA TTG TCT GGG TGT CAG AAT ATT ACT AGT ACC AAA TGC AAC TTT 293  
Ile Lys Leu Ser Gly Cys Gln Asn Ile Thr Ser Thr Lys Cys Asn Phe  
75 80 85  
TCT TCA CTC AAG CTG AAT GTT TAT GAA GAA ATT AAA TTG CGT ATA AGA 341  
Ser Ser Leu Lys Leu Asn Val Tyr Glu Glu Ile Lys Leu Arg Ile Arg  
90 95 100 105  
GCA GAA AAA GAA AAC ACT TCT TCA TGG TAT GAG GTT GAC TCA TTT ACA 389  
Ala Glu Lys Glu Asn Thr Ser Ser Trp Tyr Glu Val Asp Ser Phe Thr  
110 115 120  
CCA TTT CGC AAA GCT CAG ATT GGT CCT CCA GAA GTA CAT TTA GAA GCT 437  
Pro Phe Arg Lys Ala Gln Ile Gly Pro Pro Glu Val His Leu Glu Ala  
125 130 135

Received 2/20/2012 6:00 AM

GAA GAT AAG GCA ATA GTG ATA CAC ATC TCT CCT GGA ACA AAA GAT AGT Glu Asp Lys Ala Ile Val Ile His Ile Ser Pro Gly Thr Lys Asp Ser 140 145 150	485
GTT ATG TGG GCT TTG GAT GGT TTA AGC TTT ACA TAT AGC TTA CTT ATC Val Met Trp Ala Leu Asp Gly Leu Ser Phe Thr Tyr Ser Leu Leu Ile 155 160 165	533
TGG AAA AAC TCT TCA GGT GTA GAA GAA AGG ATT GAA AAT ATT TAT TCC Trp Lys Asn Ser Ser Gly Val Glu Glu Arg Ile Glu Asn Ile Tyr Ser 170 175 180 185	581
AGA CAT AAA ATT TAT AAA CTC TCA CCA GAG ACT ACT TAT TGT CTA AAA Arg His Lys Ile Tyr Lys Leu Ser Pro Glu Thr Thr Tyr Cys Leu Lys 190 195 200	629
GTT AAA GCA GCA CTA CTT ACG TCA TGG AAA ATT GGT GTC TAT AGT CCA Val Lys Ala Ala Leu Leu Thr Ser Trp Lys Ile Gly Val Tyr Ser Pro 205 210 215	677
GTA CAT TGT ATA AAG ACC ACA GTT GAA AAT GAA CTA CCT CCA CCA GAA Val His Cys Ile Lys Thr Thr Val Glu Asn Glu Leu Pro Pro Pro Glu 220 225 230	725
AAT ATA GAA GTC AGT GTC CAA AAT CAG AAC TAT GTT CTT AAA TGG GAT Asn Ile Glu Val Ser Val Gln Asn Gln Asn Tyr Val Leu Lys Trp Asp 235 240 245	773
TAT ACA TAT GCA AAC ATG ACC TTT CAA GTT CAG TGG CTC CAC GCC TTT Tyr Thr Tyr Ala Asn Met Thr Phe Gln Val Gln Trp Leu His Ala Phe 250 255 260 265	821
TTA AAA AGG AAT CCT GGA AAC CAT TTG TAT AAA TGG AAA CAA ATA CCT Leu Lys Arg Asn Pro Gly Asn His Leu Tyr Lys Trp Lys Gln Ile Pro 270 275 280	869
GAC TGT GAA AAT GTC AAA ACT ACC CAG TGT GTC TTT CCT CAA AAC GTT Asp Cys Glu Asn Val Lys Thr Thr Gln Cys Val Phe Pro Gln Asn Val 285 290 295	917
TTC CAA AAA GGA ATT TAC CTT CTC CGC GTA CAA GCA TCT GAT GGA AAT Phe Gln Lys Gly Ile Tyr Leu Leu Arg Val Gln Ala Ser Asp Gly Asn 300 305 310	965
AAC ACA TCT TTT TGG TCT GAA GAG ATA AAG TTT GAT ACT GAA ATA CAA Asn Thr Ser Phe Trp Ser Glu Glu Ile Lys Phe Asp Thr Glu Ile Gln 315 320 325	1013
GCT TTC CTA CTT CCT CCA GTC TTT AAC ATT AGA TCC CTT AGT GAT TCA Ala Phe Leu Leu Pro Pro Val Phe Asn Ile Arg Ser Leu Ser Asp Ser 330 335 340 345	1061
TTC CAT ATC TAT ATC GGT GCT CCA AAA CAG TCT GGA AAC ACG CCT GTG Phe His Ile Tyr Ile Gly Ala Pro Lys Gln Ser Gly Asn Thr Pro Val 350 355 360	1109
ATC CAG GAT TAT CCA CTG ATT TAT GAA ATT ATT TTT TGG GAA AAC ACT Ile Gln Asp Tyr Pro Leu Ile Tyr Glu Ile Ile Phe Trp Glu Asn Thr 365 370 375	1157
TCA AAT GCT GAG AGA AAA ATT ATC GAG AAA AAA ACT GAT GTT ACA GTT Ser Asn Ala Glu Arg Lys Ile Ile Glu Lys Lys Thr Asp Val Thr Val 380 385 390	1205
CCT AAT TTG AAA CCA CTG ACT GTA TAT TGT GTG AAA GCC AGA GCA CAC Pro Asn Leu Lys Pro Leu Thr Val Tyr Cys Val Lys Ala Arg Ala His 395 400 405	1253

ACC ATG GAT GAA AAG CTG AAT AAA AGC AGT GTT TTT AGT GAC GCT GTA	1301
Thr Met Asp Glu Lys Leu Asn Lys Ser Ser Val Phe Ser Asp Ala Val	
410 415 420 425	
TGT GAG AAA ACA AAA CCA GGA AAT ACC TCT AAA ATT TGG CTT ATA GTT	1349
Cys Glu Lys Thr Lys Pro Gly Asn Thr Ser Lys Ile Trp Leu Ile Val	
430 435 440	
GGA ATT TGT ATT GCA TTA TTT GCT CTC CCG TTT GTC ATT TAT GCT GCG	1397
Gly Ile Cys Ile Ala Leu Phe Ala Leu Pro Phe Val Ile Tyr Ala Ala	
445 450 455	
AAA GTC TTC TTG AGA TGC ATC AAT TAT GTC TTC TTT CCA TCA CTT AAA	1445
Lys Val Phe Leu Arg Cys Ile Asn Tyr Val Phe Phe Pro Ser Leu Lys	
460 465 470	
CCT TCT TCC AGT ATA GAT GAG TAT TTC TCT GAA CAG CCA TTG AAG AAT	1493
Pro Ser Ser Ser Ile Asp Glu Tyr Phe Ser Glu Gln Pro Leu Lys Asn	
475 480 485	
CTT CTG CTT TCA ACT TCT GAG GAA CAA ATC GAA AAA TGT TTC ATA ATT	1541
Leu Leu Leu Ser Thr Ser Glu Glu Gln Ile Glu Lys Cys Phe Ile Ile	
490 495 500 505	
GAA AAT ATA AGC ACA ATT GCT ACA GTA GAA GAA ACT AAT CAA ACT GAT	1589
Glu Asn Ile Ser Thr Ile Ala Thr Val Glu Glu Thr Asn Gln Thr Asp	
510 515 520	
GAA GAT CAT AAA AAA TAC AGT TCC CAA ACT AGC CAA GAT TCA GGA AAT	1637
Glu Asp His Lys Lys Tyr Ser Ser Gln Thr Ser Gln Asp Ser Gly Asn	
525 530 535	
TAT TCT AAT GAA GAT GAA AGC GAA AGT AAA ACA AGT GAA GAA CTA CAG	1685
Tyr Ser Asn Glu Asp Glu Ser Glu Ser Lys Thr Ser Glu Glu Leu Gln	
540 545 550	
CAG GAC TTT GTA TGACCAAGAAA TGAACGTGTGT CAAGTATAAG GTTTTTCAGC	1737
Gln Asp Phe Val	
555	
AGGAGTTACA CTGGTACC	1755

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 557 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met Met Val Val Leu Leu Gly Ala Thr Thr Leu Val Leu Val Ala Val	
1 5 10 15	
Gly Pro Trp Val Leu Ser Ala Ala Ala Gly Gly Lys Asn Leu Lys Ser	
20 25 30	
Pro Gln Lys Val Glu Val Asp Ile Ile Asp Asp Asn Phe Ile Leu Arg	
35 40 45	
Trp Asn Arg Ser Asp Glu Ser Val Gly Asn Val Thr Phe Ser Phe Asp	
50 55 60	

Tyr Gln Lys Thr Gly Met Asp Asn Trp Ile Lys Leu Ser Gly Cys Gln  
65 70 75 80

Asn Ile Thr Ser Thr Lys Cys Asn Phe Ser Ser Leu Lys Leu Asn Val  
85 90 95

Tyr Glu Glu Ile Lys Leu Arg Ile Arg Ala Glu Lys Glu Asn Thr Ser  
100 105 110

Ser Trp Tyr Glu Val Asp Ser Phe Thr Pro Phe Arg Lys Ala Gln Ile  
115 120 125

Gly Pro Pro Glu Val His Leu Glu Ala Glu Asp Lys Ala Ile Val Ile  
130 135 140

His Ile Ser Pro Gly Thr Lys Asp Ser Val Met Trp Ala Leu Asp Gly  
145 150 155 160

Leu Ser Phe Thr Tyr Ser Leu Leu Ile Trp Lys Asn Ser Ser Gly Val  
165 170 175

Glu Glu Arg Ile Glu Asn Ile Tyr Ser Arg His Lys Ile Tyr Lys Leu  
180 185 190

Ser Pro Glu Thr Thr Tyr Cys Leu Lys Val Lys Ala Ala Leu Leu Thr  
195 200 205

Ser Trp Lys Ile Gly Val Tyr Ser Pro Val His Cys Ile Lys Thr Thr  
210 215 220

Val Glu Asn Glu Leu Pro Pro Glu Asn Ile Glu Val Ser Val Gln  
225 230 235 240

Asn Gln Asn Tyr Val Leu Lys Trp Asp Tyr Thr Tyr Ala Asn Met Thr  
245 250 255

Phe Gln Val Gln Trp Leu His Ala Phe Leu Lys Arg Asn Pro Gly Asn  
260 265 270

His Leu Tyr Lys Trp Lys Gln Ile Pro Asp Cys Glu Asn Val Lys Thr  
275 280 285

Thr Gln Cys Val Phe Pro Gln Asn Val Phe Gln Lys Gly Ile Tyr Leu  
290 295 300

Leu Arg Val Gln Ala Ser Asp Gly Asn Asn Thr Ser Phe Trp Ser Glu  
305 310 315 320

Glu Ile Lys Phe Asp Thr Glu Ile Gln Ala Phe Leu Leu Pro Pro Val  
325 330 335

Phe Asn Ile Arg Ser Leu Ser Asp Ser Phe His Ile Tyr Ile Gly Ala  
340 345 350

Pro Lys Gln Ser Gly Asn Thr Pro Val Ile Gln Asp Tyr Pro Leu Ile  
355 360 365

Tyr Glu Ile Ile Phe Trp Glu Asn Thr Ser Asn Ala Glu Arg Lys Ile  
370 375 380

Ile Glu Lys Lys Thr Asp Val Thr Val Pro Asn Leu Lys Pro Leu Thr  
385 390 395 400

Val Tyr Cys Val Lys Ala Arg Ala His Thr Met Asp Glu Lys Leu Asn  
405 410 415

Lys Ser Ser Val Phe Ser Asp Ala Val Cys Glu Lys Thr Lys Pro Gly  
420 425 430

Asn Thr Ser Lys Ile Trp Leu Ile Val Gly Ile Cys Ile Ala Leu Phe  
435 440 445

Ala Leu Pro Phe Val Ile Tyr Ala Ala Lys Val Phe Leu Arg Cys Ile  
450 455 460

Asn Tyr Val Phe Phe Pro Ser Leu Lys Pro Ser Ser Ser Ile Asp Glu  
465 470 475 480


Tyr Phe Ser Glu Gln Pro Leu Lys Asn Leu Leu Ser Thr Ser Glu  
485 490 495

Glu Gln Ile Glu Lys Cys Phe Ile Ile Glu Asn Ile Ser Thr Ile Ala  
500 505 510

Thr Val Glu Glu Thr Asn Gln Thr Asp Glu Asp His Lys Lys Tyr Ser  
515 520 525

Ser Gln Thr Ser Gln Asp Ser Gly Asn Tyr Ser Asn Glu Asp Glu Ser  
530 535 540

Glu Ser Lys Thr Ser Glu Glu Leu Gln Gln Asp Phe Val  
545 550 555



IBM PC AT 1.44mb MSDOS 3.3  
BENOIT, et al.  
US 08/307,588  
FILED:05-DEC-1994  
"MONOCLONAL ANTIBODIES  
AGAINST THE INTERFERON..."  
ATTY DOCKET:17283/117/GUPL  
DATA REC'D:26 MAR 1996




FIGURE 1A




FIGURE 1B

**SUBSTITUTE SHEET**

CTGCAGGGATCTGGGGGGCTCCCAG

ATG ATG GTC GTC CTC CTG GGC GCG ACG ACC CTA GTG CTC GTC GCC GTG GGC CCA  
Met Met Val Val Leu Leu Gly Ala Thr Thr Leu Val Leu Val Ala Val Gly Pro

TGG GTG TTG TCC GCA GCC GCA GGT GGA AAA AAT CTA AAA TCT CCT CAA AAA GTA  
Trp Val Leu Ser Ala Ala Gly Gly Lys Asn Leu Lys Ser Pro Gln Lys Val

GAG GTC GAC ATC ATA GAT GAC AAC TTT ATC CTG AGG TGG AAC AGG AGC GAT GAG  
Glu Val Asp Ile Ile Asp Asp Asn Phe Ile Leu Arg Trp Asn Arg Ser Asp Glu

TCT GTC GGG AAT GTG ACT TTT TCA TTC GAT TAT CAA AAA ACT GGG ATG GAT AAT  
Ser Val Gly Asn Val Thr Phe Ser Asp Tyr Gln Lys Thr Gly Met Asp Asn

TGG ATA AAA TTG TCT GGG TGT CAG AAT ATT ACT AGT ACC AAA TGC AAC TTT TCT  
Trp Ile Lys Leu Ser Gly Cys Gln Asn Ile Thr Ser Thr Lys Cys Asn Phe Ser

TCA CTC AAG CTG AAT GTT TAT GAA GAA ATT AAA TTG CGT ATA AGA GCA GAA AAA  
Ser Leu Lys Leu Asn Val Tyr Glu Glu Ile Lys Leu Arg Ile Arg Ala Glu Lys

GAA AAC ACT TCT TCA TGG TAT GAG GTT GAC TCA TTT ACA CCA TTT CGC AAA GCT  
Glu Asn Thr Ser Ser Trp Tyr Glu Val Asp Ser Phe Thr Pro Phe Arg Lys Ala

CAG ATT GGT CCT CCA GAA GTA CAT TTA GAA GCT GAA GAT AAG GCA ATA GTG ATA  
Gln Ile Gly Pro Pro Glu Val His Leu Glu Ala Glu Asp Lys Ala Ile Val Ile

CAC ATC TCT CCT GGA ACA AAA GAT AGT GTT ATG TGG GCT TTG GAT GGT TTA AGC  
His Ile Ser Pro Gly Thr Lys Asp Ser Val Met Trp Ala Leu Asp Gly Leu Ser

TTT ACA TAT AGC TTA CTT ATC TGG AAA AAC TCT TCA GGT GTA GAA GAA AGG ATT  
Phe Thr Tyr Ser Leu Leu Ile Trp Lys Asn Ser Ser Gly Val Glu Arg Ile

GAA AAT ATT TAT TCC AGA CAT AAA ATT TAT AAA CTC TCA CCA GAG ACT ACT TAT  
Glu Asn Ile Tyr Ser Arg His Lys Ile Tyr Lys Leu Ser Pro Glu Thr Thr Tyr

TGT CTA AAA GTT AAA GCA GCA CTA CTT ACG TCA TGG AAA ATT GGT GTC TAT AGT  
Cys Leu Lys Val Lys Ala Ala Leu Leu Thr Ser Trp Lys Ile Gly Val Tyr Ser

CCA GTA CAT TGT ATA AAG ACC ACA GTT GAA AAT GAA CTA CCT CCA CCA GAA AAT  
Pro Val His Cys Ile Lys Thr Thr Val Glu Asn Glu Leu Pro Pro Glu Asn

ATA GAA GTC AGT GTC CAA AAT CAG AAC TAT GTT CTT AAA TGG GAT TAT ACA TAT  
Ile Glu Val Ser Val Gln Asn Gln Asn Tyr Val Leu Lys Trp Asp Tyr Thr Tyr

GCA AAC ATG ACC TTT CAA GTT CAG TGG CTC CAC GCC TTT TTA AAA AGG AAT CCT  
Ala Asn Met Thr Phe Gln Val Gln Trp Leu His Ala Phe Leu Lys Arg Asn Pro

GGA AAC CAT TTG TAT AAA TGG AAA CAA ATA CCT GAC TGT GAA AAT GTC AAA ACT  
Gly Asn His Leu Tyr Lys Trp Lys Gln Ile Pro Asp Cys Glu Asn Val Lys Thr

ACC CAG TGT GTC TTT CCT CAA AAC GTT TTC CAA AAA GGA ATT TAC CTT CTC CGC  
Thr Gln Cys Val Phe Pro Gln Asn Val Phe Gln Lys Gly Ile Tyr Leu Leu Arg

GTA CAA GCA TCT GAT GGA AAT AAC ACA TCT TTT TGG TCT GAA GAG ATA AAG TTT  
Val Gln Ala Ser Asp Gly Asn Asn Thr Ser Phe Trp Ser Glu Glu Ile Lys Phe

GAT ACT GAA ATA CAA GCT TTC CTA CTT CCT CCA GTC TTT AAC ATT AGA TCC CTT  
Asp Thr Glu Ile Gln Ala Phe Leu Leu Pro Pro Val Phe Asn Ile Arg Ser Leu

AGT GAT TCA TTC CAT ATC TAT ATC GGT GCT CCA AAA CAG TCT GGA AAC ACG CCT  
Ser Asp Ser Phe His Ile Tyr Ile Gly Ala Pro Lys Gln Ser Gly Asn Thr Pro

GTG ATC CAG GAT TAT CCA CTG ATT TAT GAA ATT ATT TTT TGG GAA AAC ACT TCA  
Val Ile Gln Asp Tyr Pro Leu Ile Tyr Glu Ile Ile Phe Trp Glu Asn Thr Ser

AAT GCT GAG AGA AAA ATT ATC GAG AAA AAA ACT GAT GTT ACA GTT CCT AAT TTG  
Asn Ala Glu Arg Lys Ile Ile Glu Lys Lys Thr Asp Val Thr Val Pro Asn Leu

AAA CCA CTG ACT GTA TAT TGT GTG AAA GCC AGA GCA CAC ACC ATG GAT GAA AAG  
Lys Pro Leu Thr Val Tyr Cys Val Ala Arg Ala His Thr Met Asp Glu Lys

CTG AAT AAA AGC AGT TTT AGT GAC GCT GTA TGT GAG AAA ACA AAA CCA GGA  
Leu Asn Lys Ser Ser Val Phe Ser Asp Ala Val Cys Glu Lys Thr Lys Pro Gly

AAT ACC TCT AAA TGA GGT ACC  
Asn Thr Ser Lys

FIGURE 2B

CTGGCAGGCACTTGCGCGCGCTCCCG

ATG ATG GTC GTC CTC CTG GGC GCG ACG ACC CTA GTG CTC GTC GCC GTG GGC CCA  
Met Met Val Val Leu Leu Ala Thr Thr Leu Val Leu Val Ala Val Gly Pro

TGG GTG TTG TCC GCA GCC GCA GGT GGA AAA AAT CTA AAA TCT CCT CAA AAA GTA  
Trp Val Leu Ser Ala Ala Gly Gly Lys Asn Leu Lys Ser Pro Gln Lys Val

GAG GTC GAC ATC ATA GAT GAC AAC TTT ATC CTG AGG TGG AAC AGC AGC GAT GAG  
Glu Val Asp Ile Ile Asp Asn Phe Ile Leu Arg Trp Asn Arg Ser Asp Glu

TCT GTC GGG AAT GTG ACT TTT TCA TTC GAT TAT CAA AAA ACT GGG ATG GAT AAT  
Ser Val Gly Asn Val Thr Phe Ser Phe Asp Tyr Gln Lys Thr Gly Met Asp Asn

TGG ATA AAA TTG TCT GGG TGT CAG AAT ATT ACT AGT ACC AAA TGC AAC TTT TCT  
Trp Ile Lys Leu Ser Gly Cys Gln Asn Ile Thr Ser Thr Lys Cys Asn Phe Ser

TCA CTC AAG CTG AAT GTT TAT GAA GAA ATT AAA TTG CGT ATA AGA GCA GAA AAA  
Ser Leu Lys Leu Asn Val Tyr Glu Glu Ile Lys Leu Arg Ile Arg Ala Glu Lys

GAA AAC ACT TCT TCA TGG TAT GAG GTT GAC TCA TTT ACA CCA TTT CGC AAA GCT  
Glu Asn Thr Ser Ser Trp Tyr Glu Val Asp Ser Phe Thr Pro Phe Arg Lys Ala

CAG ATT GGT CCT CCA GAA GTA CAT TTA GAA GCT GAA GAT AAG GCA ATA GTG ATA  
Gln Ile Gly Pro Pro Glu Val His Leu Glu Ala Glu Asp Lys Ala Ile Val Ile

CAC ATC TCT CCT GGA ACA AAA GAT AGT GTT ATG TGG GCT TTG GAT GGT TTA AGC  
His Ile Ser Pro Gly Thr Lys Asp Ser Val Met Trp Ala Leu Asp Gly Leu Ser

TTT ACA TAT AGC TTA CTT ATC TGG AAA AAC TCT TCA GGT GTA GAA GAA AGG ATT  
Phe Thr Tyr Ser Leu Leu Ile Trp Lys Asn Ser Ser Gly Val Glu Glu Arg Ile

GAA AAT ATT TAT TCC AGA CAT AAA ATT TAT AAA CTC TCA CCA GAG ACT ACT TAT  
Glu Asn Ile Tyr Ser Arg His Lys Ile Tyr Lys Leu Ser Pro Glu Thr Thr Tyr

TGT CTA AAA GTT AAA GCA GCA CTA CTT ACG TCA TGG AAA ATT GGT GTC TAT AGT  
Cys Leu Lys Val Lys Ala Ala Leu Leu Thr Ser Trp Lys Ile Gly Val Tyr Ser

CCA GTA CAT TGT ATA AAG ACC ACA GTT GAA AAT GAA CTA CCT CCA CCA GAA AAT  
Pro Val His Cys Ile Lys Thr Thr Val Glu Asn Glu Leu Pro Pro Pro Glu Asn

ATA GAA GTC AGT GTC CAA AAT CAG AAC TAT GTT CTT AAA TGG GAT TAT ACA TAT  
Ile Glu Val Ser Val Gln Asn Gln Asn Tyr Val Leu Lys Trp Asp Tyr Thr Tyr

GCA AAC ATG ACC TTT CAA GTT CAG TGG CTC CAC GCC TTT TTA AAA AGG AAT CCT  
Ala Asn Met Thr Phe Gln Val Gln Trp Leu His Ala Phe Leu Lys Arg Asn Pro

GGA AAC CAT TTG TAT AAA TGG AAA CAA ATA CCT GAC TGT GAA AAT GTC AAA ACT  
Gly Asn His Leu Tyr Lys Trp Lys Gln Ile Pro Asp Cys Glu Asn Val Lys Thr

ACC CAG TGT GTC TTT CCT CAA AAC GTT TTC CAA AAA GGA ATT TAC CTT CTC CGC  
Thr Gln Cys Val Phe Pro Gln Asn Val Phe Gln Lys Gly Ile Tyr Leu Leu Arg

GTA CAA GCA TCT GAT GGA AAT AAC ACA TCT TTT TGG TCT GAA GAG ATA AAG TTT  
Val Gln Ala Ser Asp Gly Asn Asn Thr Ser Phe Trp Ser Glu Glu Ile Lys Phe

GAT ACT GAA ATA CAA GCT TTC CTA CTT CCT CCA GTC TTT AAC ATT AGA TCC CTT  
Asp Thr Glu Ile Gln Ala Phe Leu Leu Pro Pro Val Phe Asn Ile Arg Ser Leu

AGT GAT TCA TTC CAT ATC TAT ATC GGT GCT CCA AAA CAC TGT GGA AAG AGC CCT  
 Ser Asp Ser Phe His Ile Tyr Ile Gly Ala Pro Lys Gln Ser Gly Asn Thr Pro

G TG ATC CAG GAT TAT CCA CTG ATT TAT GAA ATT ATT TTT TGG GAA AAC ACT TCA  
 Val Ile Gln Asp Tyr Pro Leu Ile Tyr Glu Ile Phe Trp Glu Asn Thr Ser

AAT GCT GAG AGA AAA ATT ATC GAG AAA AAA ACT GAT GTT ACA GTT CCT AAT TTG  
 Asn Ala Glu Arg Lys Ile Ile Glu Lys Thr Asp Val Thr Val Pro Asn Leu

AAA CCA CTG ACT GTA TAT TGT GTG AAA GCC AGA GCA CAC ACC ATG GAT GAA AAG  
 Lys Pro Leu Thr Val Tyr Cys Val Lys Ala Arg Ala His Thr Met Asp Glu Lys

CTG AAT AAA AGC AGT GTT TTT AGT GAC GCT GTA TGT GAG AAA ACA AAA CCA GGA  
 Leu Asn Lys Ser Ser Val Phe Ser Asp Ala Val Cys Glu Lys Thr Lys Pro Gly

AAT ACC TCT AAA ATT TGG CTT ATA GTT GGA ATT TGT ATT GCA TTA TTT GCT CTC  
 Asn Thr Ser Lys Ile Trp Leu Ile Val Gly Ile Cys Ile Ala Leu Phe Ala Leu

CCG TTT GTC ATT TAT GCT GCG AAA GTC TTC TTG AGA TGC ATC AAT TAT GTC TTC  
 Pro Phe Val Ile Tyr Ala Ala Lys Val Phe Leu Arg Cys Ile Asn Tyr Val Phe

TTT CCA TCA CTT AAA CCT TCT TCC AGT ATA GAT GAG TAT TTC TCT GAA CAG CCA  
 Phe Pro Ser Leu Lys Pro Ser Ser Ile Asp Glu Tyr Phe Ser Glu Gln Pro

TTG AAG AAT CTT CTG CTT TCA ACT TCT GAG GAA CAA ATC GAA AAA TGT TTC ATA  
 Leu Lys Asn Leu Leu Ser Thr Ser Glu Glu Gln Ile Glu Lys Cys Phe Ile

ATT GAA AAT ATA AGC ACA ATT GCT ACA GTA GAA GAA ACT AAT CAA ACT GAT GAA  
 Ile Glu Asn Ile Ser Thr Ile Ala Thr Val Glu Glu Thr Asn Gln Thr Asp Glu

GAT CAT AAA AAA TAC AGT TCC CAA ACT AGC CAA GAT TCA GGA AAT TAT TCT AAT  
 Asp His Lys Lys Tyr Ser Ser Gln Thr Ser Gln Asp Ser Gly Asn Tyr Ser Asn

GAA GAT GAA AGC GAA AGT AAA ACA AGT GAA GAA CTA CAG CAG GAC TTT GTA TGA  
 Glu Asp Glu Ser Glu Ser Lys Thr Ser Glu Glu Leu Gln Gln Asp Phe Val

CCAGAAATGAACGTGTCAAGTATAAGGTTTCAGCAGGAGTTACACTGGTACC

FIGURE 3B

SUBSTITUTE SHEET

SEARCHED  
INDEXED  
MAILED  
SERIALIZED  
FILED

# DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address, and citizenship are as stated below next to my name. I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**MONOCLONAL ANTIBODIES AGAINST THE INTERFERON RECEPTOR, WITH NEUTRALIZING ACTIVITY AGAINST TYPE 1 INTERFERON**

the specification of which (check one)

is attached hereto

was filed on March 30, 1993 as Application Serial No. PCT/EP93/00770 and was amended on \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is known by me to be material to patentability as defined in Title 37, Code of Federal Regulations § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

**PRIOR FOREIGN APPLICATION(S)**

NUMBER	COUNTRY	DAY/MONTH/YEAR FILED	PRIORITY CLAIMED
<u>92400902.0</u>	<u>European</u>	<u>31/March/1992</u>	<u>Yes</u>


I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) listed below and insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is known by me to be material to patentability as defined in Title 37, Code of Federal Regulations § 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

APPLICATION SERIAL NO.	FILING DATE	STATUS: PATENTED, PENDING, ABANDONED

I hereby appoint as my attorneys, with full powers of substitution and revocation, to prosecute this application and transact all business in the Patent and Trademark Office connected therewith: Stephen A. Bent, Reg. No. 29,768; David A. Blumenthal, Reg. No. 26,257; John J. Feldhaus, Reg. No. 28,822; Donald D. Jeffery, Reg. No. 19,980; Peter G. Mack, Reg. No. 26,001; Brian J. McNamara, Reg. No. 32,789; Sybil Meloy, Reg. No. 22,749; George E. Quillin, Reg. No. 32,792; Colin G. Sandercock, Reg. No. 31,298; Bernhard D. Saxe, Reg. No. 28,665; Richard L. Schwaab, Reg. No. 25,479; Arthur Schwartz, Reg. No. 22,115.

Send all correspondence to **FOLEY & LARDNER**, 3000 K Street, N.W., Suite 500, P.O. Box 25696, Washington, D.C. 20007-8696. Address telephone communications to Bernhard D. Saxe at (202) 672-5300.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.


Full Name of First or Sole Inventor <i>Patrick BENOIT</i>	Signature of First or Sole Inventor 	Date <i>14 Nov/94</i>
Residence Address <i>Paris, France</i>	Country of Citizenship <i>FRANCE</i>	
Post Office Address <i>24, rue Jonquoy, F-75014 Paris, France</i>		

Signatures should conform to names as typewritten.  Additional inventors on attached Page 2.

Full Name of Second Inventor <i>Francois MEYER</i>	Signature of Second Inventor 	Date
Residence Address <i>Paris, France</i>	Country of Citizenship <b>FRANCE</b>	
Post Office Address <i>14, square Adanson, F-75005 Paris, France</i>		

Full Name of Third Inventor <i>Debborah MAGUIRE</i>	Signature of Third Inventor 	Date
Residence Address <i>Paris, France</i>	Country of Citizenship <b>FRANCE</b>	
Post Office Address <i>24, rue Maitre-Albert, F-75005 Paris, France</i>		

Full Name of Fourth Inventor <i>Ivan PLAVEC</i>	Signature of Fourth Inventor 	Date
Residence Address <i>Fresnes, France</i>	Country of Citizenship <b>FRANCE</b>	
Post Office Address <i>1, allee du Capitaine-Dupont F-92260 Fresnes, France</i>		



Full Name of Fifth Inventor <i>Michael G. TOVEY</i>	Signature of Fifth Inventor 	Date <u>11/11/94</u>
Residence Address <i>Paris, France</i>	Country of Citizenship <b>GREAT BRITAIN</b>	
Post Office Address <i>6, rue des Quatrefages, F-75005 Paris, France</i>		



SEARCHED INDEXED SERIALIZED FILED



PAGE 2



Docket No. 17283/117/GUPL

B 1608AC

Full Name of Second Inventor <b>Francois MEYER</b>	Signature of Second Inventor 	Date 
Residence Address <b>Paris, France</b>	Country of Citizenship <b>FRANCE</b> <b>LUXEMBOURG</b>	
Post Office Address <b>14, square Adanson, F-75005 Paris, France</b>		

Full Name of Third Inventor <b>Deborah MAGUIRE</b>	Signature of Third Inventor 	Date 
Residence Address <b>Paris, France</b>	Country of Citizenship <b>FRANCE</b> <b>UNITED KINGDOM</b>	
Post Office Address <b>24, rue Maitre-Albert, F-75005 Paris, France</b>		

Full Name of Fourth Inventor <b>Ivan PLAVEC</b>	Signature of Fourth Inventor 	Date 
Residence Address <b>Fresnes, France</b>	Country of Citizenship <b>FRANCE</b> <b>CROATIA</b>	
Post Office Address <b>1, allee du Capitaine-Dupont F-92260 Fresnes, France</b>		

Full Name of Fifth Inventor <b>Michael G. TOVEY</b>	Signature of Fifth Inventor 	Date 
Residence Address <b>Paris, France</b>	Country of Citizenship <b>GREAT BRITAIN</b>	
Post Office Address <b>6, rue des Quatreages, F-75005 Paris, France</b>		

B 1608 AC

PAGE 2

Docket No. 17283/117/GUPL

Full Name of Second Inventor <i>Francois MEYER</i>	Signature of Second Inventor <i>X. Meyer</i>	Date <i>11/11/94</i>
Residence Address <i>Paris, France</i>	Country of Citizenship <i>FRANCE L'AYETBOURG</i>	
Post Office Address <i>14, square Adanson, F-75005 Paris, France</i>		

Full Name of Third Inventor <i>Deborah MAGUIRE</i>	Signature of Third Inventor <i>D. Maguire</i>	Date
Residence Address <i>Paris, France</i>	Country of Citizenship <i>FRANCE</i>	
Post Office Address <i>24, rue Maitre-Albert, F-75005 Paris, France</i>		

Full Name of Fourth Inventor <i>Ivan PLAVEC</i>	Signature of Fourth Inventor <i>Ivan Plavec</i>	Date <i>11/11/94</i>
Residence Address <i>Fresnes, France</i>	Country of Citizenship <i>FRANCE CROATIA</i>	
Post Office Address <i>1, allee du Capitaine-Dupont F-92260 Fresnes, France</i>		

Full Name of Fifth Inventor <i>Michael G. TOVEY</i>	Signature of Fifth Inventor <i>Michael Tovey</i>	Date
Residence Address <i>Paris, France</i>	Country of Citizenship <i>GREAT BRITAIN</i>	
Post Office Address <i>6, rue des Quatre fages, F-75005 Paris, France</i>		